

Raffinose Oligosaccharide Catabolism in *Arabidopsis thaliana*

Dissertation

zur

**Erlangung der naturwissenschaftlichen Doktorwürde
(Dr. sc. nat.)**

vorgelegt der

Mathematisch-naturwissenschaftlichen Fakultät

der

Universität Zürich

von

Aurélie Egert

aus

Frankreich

Promotionskomitee

Prof. Dr. Enrico Martinoia (Vorsitz)

Prof. Dr. Felix Keller (Leitung der Dissertation)

Dr. Norbert Sprenger

Zürich, 2011

INDEX

SUMMARY	v
ZUSAMMENFASSUNG	vii
CHAPTER I: General Introduction	1
1.1 RFO biosynthesis	5
1.1.1 Galactinol synthase (GolS) catalyzes the formation of the RFO precursor Gol.....	6
1.1.2 Raffinose synthase (RafS) catalyzes the synthesis of the first RFO member	7
1.1.3 Stachyose synthase (StaS) catalyzes the synthesis of the second RFO member	8
1.1.4 Higher DP RFO are mainly synthesized <i>via</i> a Gol-independent pathway.....	9
1.2 RFO catabolism	11
1.2.1 RFO degradation starts with the hydrolytic action of α -D-galactosidases (α Gals)	12
1.2.2 Galactokinase (GALK) phosphorylates free Gal produced by the degradation of Gal-containing compounds	16
1.2.3 Free Gal is toxic.....	17
1.2.4 The Gal salvage pathway continues	18
1.3 RFOs play a role in abiotic stress tolerance.....	21
1.3.1 Plants are frequently exposed to abiotic stress.....	21
1.3.2 RFOs accumulate during stress and may act as osmoprotectants.....	23
1.3.3 Cold stress involves molecular regulation and incudes RFO accumulation	25
1.3.4 Water deficit and salt stress may induce RFO accumulation.....	28
1.3.5 Other abiotic stress and RFO accumulation	29
1.4 RFOs play a role in carbon partitioning and storage.....	31
1.4.1 Transport RFOs move in the phloem from source to sink	31
1.4.2 Storage RFOs accumulate in the vacuole and can be easily mobilized.....	33
1.5 Aims of my thesis	35
CHAPTER II: Functional identification of Arabidopsis alkaline α-galactosidases ATSIP1 (At1g55740) and ATSIP2 (At3g57520)	37
2.1 Introduction	39
2.2 Materials and methods.....	41
2.2.1 Plants material and growth conditions	41
2.2.2 Screening for T-DNA insertion lines.....	41
2.2.3 Heterologous expression of ATSIP1 and -2 in <i>Sf9</i> insect cells.....	42
2.2.3.1 Bacmid construction and transfection	42
2.2.3.2 Expression of ATSIP1 and -2 in <i>Sf9</i> cells	43
2.2.3.3 RNA isolation from <i>Sf9</i> cells and cDNA synthesis.....	44
2.2.4 α -Galactosidase activity assays of recombinant ATSIP1 and -2.....	45
2.2.5 Desalting of extracts.....	45
2.2.6 HPLC-PAD analysis.....	46
2.2.7 Biochemical characterization of recombinant ATSIP1 and -2.....	46

2.2.8 Cold acclimation and deacclimation treatments	47
2.2.9 Water-deficit and rehydration treatments	47
2.2.10 Plant enzyme extraction and activity assays	47
2.2.11 EDTA-mediated phloem exudation and ¹⁴ C pulse-chase experiment.....	48
2.2.12 RNA isolation from plant tissue and semi-quantitative PCR (sqPCR).....	48
2.2.13 Promoter-β-glucuronidase (GUS) plants	49
2.2.14 Histochemical staining for β-glucuronidase (GUS) activity	49
2.2.15 Preparation of root cross sections for light microscopy	50
2.2.16 Seed germination assays	50
2.2.17 Water soluble carbohydrate (WSC) extractions.....	50
2.3 Results.....	51
2.3.1 ATSIP1 and -2 are <i>bona-fide</i> αGals with different substrate specificities	51
2.3.2 Biochemical characterization of recombinant ATSIP1 and -2	53
2.3.3 ATSIP2 does not seem to play an important role during deacclimation of cold-acclimated plants	54
2.3.4 ATSIP2 does not seem to play an important role during rehydration of dehydrated plants...	56
2.3.5 Investigating the role of ATSIP1 and -2 role in phloem unloading in leaves	58
2.3.5.1 ATSIP2 has a sink-specific expression and αGal activity pattern	60
2.3.5.2 ATSIP1 has a sink-specific expression pattern but the Sta and Gol-specific αGal activity remains low	61
2.3.6 Physiological functions of ATSIP1 and -2 during seed germination	62
2.4 Discussion	65
CHAPTER III: AtDIN10 (At5g20250) is an alkaline αGal located in the chloroplast.....	67
3.1 Introduction	69
3.2 Materials and methods.....	71
3.2.1 Plant material and growth conditions	71
3.2.2 Recombinant AtDIN10 expression and characterization in <i>Sf9</i> insect cells	71
3.2.3 GFP fusion protein analysis	71
3.2.4 Biochemical characterization of recombinant AtDIN10	72
3.3 Results.....	73
3.3.1 <i>AtDIN10</i> contains a putative chloroplast transit peptide (cTP).....	73
3.3.2 AtDIN10 is localized in the chloroplast.....	73
3.3.3 AtDIN10 is an alkaline αGal	74
3.4 Discussion	76
CHAPTER IV: An Arabidopsis T-DNA insertion mutant for galactokinase (AtGALK, At3g06580) hyperaccumulates free galactose and exhibits insensitivity to exogenous galactose.....	79
4.1 Introduction.....	81
4.2 Materials and methods.....	83
4.2.1 Plant materials and growth conditions	83

4.2.2	Screening for T-DNA insertion lines.....	83
4.2.3	Recombinant AtGALK expression in <i>Sf9</i> insect cells.....	83
4.2.4	GALK activity assay and biochemical characterization of recombinant AtGALK.....	84
4.2.5	GALK analysis of plant crude extract	84
4.2.6	Construction of <i>atgalk/35S: AtGALK</i> rescued lines	84
4.2.7	RNA isolation and semi-quantitative PCR (sqPCR)	84
4.2.8	Water soluble carbohydrate (WSCs) extraction.....	85
4.2.9	Isolation of mesophyll protoplasts and vacuoles from Arabidopsis	85
4.2.10	Enzymatic quantification of Gal.....	85
4.2.11	Gal toxicity assays.....	86
4.3	Results	87
4.3.1	AtGALK was functionally expressed and biochemically characterized	87
4.3.2	Characterization of an <i>atgalk</i> T-DNA insertion line	88
4.3.3	Functional rescue of <i>atgalk</i>	90
4.3.4	<i>In vitro</i> -grown <i>atgalk</i> seedlings exhibit insensitivity to exogenous Gal	91
4.3.5	Free Gal accumulates in leaf mesophyll vacuoles of <i>atgalk</i> plants.....	92
4.3.6	Uptake mechanisms for free Gal are unaffected in <i>atgalk</i> plants	93
4.4	Discussion.....	96
CHAPTER V:	General summary, conclusions and outlook	101
5.1	General discussions and conclusions	103
5.1.1	AtSIP2 is a <i>bona fide</i> Raf-specific alkaline α Gal proposed to be involved in phloem unloading	105
5.1.2	ATSIP1 is a <i>bona fide</i> Sta- and Gol-specific alkaline α Gal with a yet unknown physiological function.....	106
5.1.3	AtDIN10 is an alkaline α Gal located in the chloroplast, proposed to degrade DGDG during dark-induced senescence	107
5.1.4	Are the other annotated RafSs genuine RafSs?.....	108
5.1.5	AtGALK is a <i>bona fide</i> GALK and the study of <i>atgalk</i> reveals a putative new Gal detoxification pathway involving a tonoplast transporter	109
5.1.6	Does the conversion of β -form of Gal to α -form via a mutarotase exist in Arabidopsis?.....	111
5.2	Outlook.....	114
REFERENCE LIST.....		115
APENDICES		
I	List of abbreviations.....	129
II	PETERS S., EGERT A., STIEGER B., KELLER F. (2010) Plant and Cell Physiology	131
III	<i>Curriculum Vitae</i>	137
IV	Acknowledgments	139