

On the Chromosomal Architecture of *Arabidopsis thaliana*

Dissertation

zur

Erlangung der naturwissenschaftlichen Doktorwürde
(Dr. sc. nat.)

vorgelegt der

Mathematisch-naturwissenschaftlichen Fakultät

der

Universität Zürich

von

Stefan Grob

von

Wallisellen/ZH und Dinhard/ZH

Promotionskomitee

Prof. Dr. Ueli Grossniklaus (Vorsitz und Leitung der Dissertation)

Dr. Thomas Wicker

Prof. Dr. Nathan Luedtke

Prof. Dr. Dirk Schübeler

Zürich, 2014

Table of Contents

Abstract	3
Zusammenfassung	4
Curriculum Vitae	5
Table of Contents.....	6
General Introduction.....	9
The Discovery of the Nucleus as Carrier of Genetic Material	9
3C Technology Permits to Study Chromosomal Architecture in High Resolution	12
Structural Components of the Nucleus	17
<i>The Nuclear Membrane</i>	17
<i>The Nuclear Matrix</i>	22
Interphase Chromosomes.....	24
Chromosomal Architecture Is Tightly Coupled to the Epigenetic Landscape	26
References General Introduction	30
Chapter I: Characterization of Chromosomal Architecture in <i>Arabidopsis</i> by Chromosome Conformation Capture	37
Chapter II: HiC Analysis in <i>Arabidopsis</i> Identifies the <i>KNOT</i>, a Structure with Similarities to the <i>flamenco</i> Locus of <i>Drosophila</i>	76
Summary.....	77
Highlights	78
Introduction	79
RESULTS.....	82
<i>Chromosomal Neighborhood</i>	82
<i>HiC Interactions Form Defined Interaction Domains</i>	82
<i>Principal Component Analysis Reveals Distinct Chromatin States</i>	83
<i>Open and Closed Chromatin Correlate with Epigenetic Chromatin States</i>	85
<i>Arabidopsis Mutants Affecting Nuclear Size Affect the Interactome</i>	88
<i>Differences between <i>crwn1</i>, <i>crwn4</i> and <i>Col-0</i> Cluster in Defined Domains</i>	89
<i>Domain Organization of Chromosome Arms Does not Change in <i>crwn1</i> and <i>crwn4</i> Mutants</i>	93
<i>Distance-dependent Decay of Interactions</i>	93
<i>Specific Chromosome Interactions Form the KNOT</i>	96
<i>FISH Confirms the Existence of the KNOT</i>	96
<i>KEEs Share Common Sequence Motifs</i>	100
<i>KEEs Show a Specific Enrichment of Epigenetic and Genomic Features</i>	101
<i>KEEs Are Preferred Transposable Element Insertions Sites</i>	102
DISCUSSION	106
<i>There Is no Distinct Chromosomal Neighbourhood for a Given Chromosome</i>	106
<i>Arabidopsis Chromosomes Show a Simple Organization with Respect to their Epigenetic Landscape and Interactome</i>	106
<i>Nuclear Morphology Affects trans-chromosomal Interactions but not Domain Structure in Arabidopsis Nuclei</i>	107
<i>Stochastic Variability between Interactomes Has to Be Carefully Assessed to Draw Biologically Relevant Conclusions</i>	108
<i>Interaction Decay Exponents Indicate a Distinct Chromatin Organization of Chromatin Arms and Pericentromeric Repeats</i>	109

<i>The KNOT Plays a Role as a TE Trap Similar to the flamenco Locus in Drosophila</i>	111
Experimental Procedures	114
Plant Material	114
HiC Experiments	114
FISH Experiments	114
Data Analysis	114
Accession Numbers	114
Author Contributions	115
Acknowledgements	115
Supplemental Information	116
Supplementary Tables	116
Extended Experimental Procedures	122
Plant Material	122
Fluorescence in situ Hybridization (FISH)	122
HiC Sample Preparation	125
HiC Library Preparation	128
HiC Sequencing Data Processing	129
Data on Epigenetic and Genomic Features	130
Calculation of the Interaction Frequency Decay Exponent	130
Determination of Chromosomal Neighborhoods	131
Identification of Chromatin Domains	131
Epigenetic Landscape and Chromatin Domains	132
Identification of KEE Locations	132
Random Sampling Strategy for Analysis of KEE and KEE Homologous Regions	133
Enrichment of Interaction Frequencies between KEE and KEE Homologous Regions	133
Enrichment of Epigenetic or Genomic Features in KEE Regions	134
Epigenetic Variance among KEE Regions	134
Occurrence of Natural Transposon Insertions in KEE Regions	134
Difference Between HiC Data Sets	135
Interaction Frequencies of Drosophila piRNA Clusters	136
References Chapter II	138
Chapter III: Additional Analyses of HiC Interactomes of Arabidopsis	142
Summary	143
Distorted Distributions of Interaction Frequencies Greatly Contribute to Observed Differences between HiC Interactomes	143
Introduction	143
Results	143
Discussion	149
Re-evaluation of the Effects of <i>morc6-1</i> on Chromosomal Architecture	151
Introduction	151
Results	152
Discussion	161
Materials and Methods	162
Distal Positions Exhibit Increased <i>Inter</i> -Chromosomal Interactions	163
Introduction	163
Results	163
Discussion	167
Methods	168
<i>Inter</i> -Chromosomal Interactome	169

<i>Introduction</i>	169
<i>Results</i>	169
<i>Discussion</i>	175
<i>Methods</i>	178
Transgenes Potentially Influence Chromosomal Architecture	180
<i>Introduction</i>	180
<i>Results</i>	180
<i>Discussion</i>	187
References Chapter III	188
General Discussion	189
3C Technologies Greatly Aid to the Understanding of Chromosomal Architecture	189
Topological Chromatin Domains	194
The KNOT and the Transgenes	196
<i>The KNOT</i>	196
<i>The Transgenes</i>	198
References General Discussion	200
Appendix: <i>Trans</i>-generational Epigenetic Inheritance of Heat Stress Response	202
<i>Introduction</i>	202
General experimental setup	203
Experimental Work-Flow	204
Results	207
<i>Growth in Heat Stress Exposed Plants is Impaired</i>	207
<i>Growth Rates Are Independent of Growth Rates of Previous Generation</i>	207
<i>Phenotypic Variation in Heat Stress Tolerance Does not Increase Over Subsequent Generations</i>	211
References Appendix	213
Acknowledgments	214